

Search Authors...



Translate Website

Search...



GR Newsletter, Enter Email



US Nato War Economy Civil Rights Environment Poverty Media Justice 9/11 War Crimes Militarization History Science

## ARNm detectat în laptele matern după vaccinarea împotriva COVID-19 poate fi transmis sugarilor: un nou studiu

Un nou studiu a constatat că femeile care alăptează vaccinate recent cu un vaccin ARNm aveau ARNm detectabil în laptele matern care se poate transfera la copilul lor.

De [Megan Redshaw](#)

Tema: Știință și Medicină

Cercetare globală, 09 octombrie 2023

[The Epoch Times](#) 27 septembrie 2023

Translate Website



Toate articolele Global Research pot fi citite în 51 de limbi prin activarea butonului Traducere site-ul web de sub numele autorului.

Pentru a primi Newsletter-ul zilnic al Global Research (articole selectate), [faceți clic aici](#).

Faceți clic pe butonul de distribuie de mai sus pentru a trimite prin e-mail/redirecționare acest articol prietenilor și colegilor dvs. Urmărește-ne pe [Instagram](#) și [Twitter](#) și abonează-te la [canalul nostru Telegram](#). Simțiți-vă liber să repostați și să distribuiți pe scară largă articolele Global Research.

\*\*\*

*Un nou studiu Lancet se adaugă la un număr tot mai mare de dovezi care sugerează că ARNm din vaccinurile COVID-19 nu rămâne la locul injectării, ci „se răspândește sistemic” în tot organismul și poate ajunge chiar în laptele matern transmis sugarilor de către cei vaccinați. mamelor.*

### Studiul a găsit ARNm în 70% din probele de lapte matern

În lucrarea Lancet [publicată în septembrie](#), cercetătorii au detectat ARNm la 70 la sută dintre femeile care au furnizat mostre de lapte matern până la 45 de ore după vaccinare. Deși ARNm detectat a fost în mare parte fragmentat și a păstrat doar 12 până la 25% din integritatea sa originală, cercetătorii au spus că sunt necesare studii suplimentare pentru a determina cantitatea minimă de ARNm care ar putea declanșa un răspuns imun la nou-născuți.



Captură de ecran de la The Lancet

Cercetătorii au colectat mostre de lapte matern de la 13 femei sănătoase, care alăptează, înainte de vaccinare și de cel puțin două ori pe zi timp de cinci zile după vaccinare. Șapte mame au furnizat mostre de lapte matern după ce au primit prima și a doua doză de vaccin, rezultând în total 20 de expuneri la vaccin și 154 de mostre de lapte matern.

Vaccine mRNA was then examined in whole breast milk and breast milk extracellular vesicles. Extracellular vesicles (EVs) are small delivery vehicles released by cells that carry biomolecules such as protein, lipids, noncoding RNA, microRNA, messenger RNA, and DNA. Breast milk contains large amounts of EVs that regulate gene expression, immune function, development, and infant growth.

Of the 13 lactating women who received either the Moderna or Pfizer vaccine, trace mRNA amounts were detected in 10 of 20 exposures up to 45 hours following vaccination. According to the study, the SARS-CoV-2 spike protein was not expressed.

Before breast milk collections, all participants tested negative for COVID-19, and prevaccination samples were negative for COVID-19 vaccine mRNA. Although all women were instructed to provide samples of 5 milliliters or greater, actual amounts were often below the threshold, hindering the experiment.

“Our proposed model suggests that after intramuscular administration, the vaccine mRNA enclosed in lipid nanoparticles is transported to the mammary glands through either hematogenous or lymphatic pathways,” the researchers wrote. “Within the mammary cells’ cytosol, a portion of the released vaccine mRNA is recruited and packaged into the developing extracellular vesicles, which are then released into the breast milk.”

## Safety Assessment of Future mRNA-Based Therapies Needed

The researchers said the significance of their research extends beyond the scope of mRNA COVID-19 vaccines and offers “valuable insights into the transport and presence of vaccine mRNA in breast milk, which can be relevant for assessing the safety and efficacy of future mRNA-based therapies administered to lactating women.”

Lead author Dr. Nazeeh Hanna, chief of the division of neonatology at NYU Langone Hospital and professor of pediatrics at NYU Long Island School of Medicine, told The Epoch Times in an email that it’s unfortunate so many people have overlooked the real value of the study.

“It is not about the COVID vaccine only but future mRNA therapy that is currently being developed,” Dr. Hanna said. “We need to make sure we get it right for the next time. This is what we wrote in the conclusion of the Lancet paper.”

Based on the study’s results, Dr. Hanna said he doesn’t believe there’s a risk of breastfeeding after receiving the vaccine if a mother waits two days following vaccination, as they did not detect vaccine mRNA after that time. He also does not feel there’s a risk if a mother is to get vaccinated and is nursing a child older than 6 months of age because the Centers for Disease Control and Prevention (CDC) allows for the vaccination of infants at 6 months of age or older.

However, mothers with infants under 6 months of age should consult their doctor, who will propose one of two options, according to Dr. Hanna:

“Continue breastfeeding since the associated risk is low,” or “temporarily pause breastfeeding for these two days post-vaccination and use previously stored breast milk to feed the child—this is a common practice by many breastfeeding mothers. Once the 2-day period is over, the mother can resume breastfeeding.”

## Other Studies Suggest mRNA Can Transfer Through Breast Milk

Pfizer and Moderna [excluded pregnant and lactating women](#) from their clinical trials and never conducted human pharmacokinetic studies with COVID-19 vaccines. Pharmacokinetic studies assess how the human body interacts with mRNA (also referred to as “[modified RNA](#)”) in vaccines for the entire duration of exposure, showing how the contents in vaccines travel throughout the body and how they’re excreted.

Despite the unknowns, the CDC and the American College of Obstetrics and Gynecology [began recommending](#) COVID-19 vaccines for pregnant and lactating women in July 2021.

The Lancet study is not the first to suggest that mRNA from COVID-19 vaccines can be potentially transferred to infants through breast milk in recently vaccinated mothers. An August 2021 study [published in Nature](#) found that lactating mothers who received Pfizer’s COVID-19 vaccine secreted SARS-CoV-2-specific IgA and IgG antibodies into breast milk, with the most significant increase at three to seven days after the second vaccine dose.

A 2022 study in [Frontiers Immunology](#) analyzed samples from 35 breastfeeding mothers and found that most mothers had “detectable SARS-CoV-2 antibody isotypes and neutralizing antibodies in serum and breastmilk,” especially after their second Pfizer vaccine dose. Although researchers concluded a “likely lack of significant exposure or sensitization” of breastmilk-fed infants of vaccinated mothers from low levels of mRNA present in breast milk, they only analyzed the serums of five infants over 8 months of age to reach that conclusion.

## Adverse Events Reported in Nursing Infants

According to the CDC, “[available data](#) on the safety of COVID-19 vaccination while breastfeeding indicates no severe reactions after the first or second dose, neither in the breastfeeding person nor the breastfed child,” and there has been “no evidence to suggest that COVID-19 vaccines are harmful to either people who have received a vaccine and are breastfeeding or to their babies.”

Yet confidential Pfizer documents obtained through a Freedom of Information Act lawsuit in April 2021—before the June 2021 recommendation that pregnant and lactating women get vaccinated—[provide evidence](#) for 215 reports of infant exposure to Pfizer’s COVID-19 vaccines through lactation reported to the company’s safety database.

Of the 215 reports, 174 were reports of “exposure via breastmilk/maternal exposure during breastfeeding.” The other 41 cases listed adverse events in infants indirectly exposed to Pfizer’s COVID-19 vaccine through breastmilk, such as fever, facial paralysis, lymphadenopathy, and vomiting.

A 2021 [data analysis](#) from the CDC’s Vaccine Adverse Event Reporting System conducted by immunologist and molecular biologist Jessica Rose, who holds a doctorate in computational biology, showed 177 adverse events. Three reports were related to infants under age one exposed through maternal vaccination.

“VAERS data confirms in recently reported results that breastfeeding infants of mothers recently injected with the COVID-19 products are succumbing to adverse events ranging from high fevers (104) lasting for days to hypophagia (undereating),” Ms. Rose told The Epoch Times in an email.

“The following is the symptom text reported for an infant (VAERS ID: 1124474) who was exposed via breast milk at the age of 12 months. ‘Mother of 12-month-old boy received the first dose of COVID-19 vaccine at 9:15 am she breastfed her 12-month-old son 3 hours later, and while breastfeeding, the child developed acute anaphylaxis,’” Ms. Rose said. “To be clear: mother had the vaccine, and the child had the reaction.”

A 2021 study [published in Breastfeeding Medicine](#) found that more than 85 percent of 180 breastfeeding women who received an mRNA COVID-19 vaccine reported local or systemic symptoms, with higher frequency following the second dose. “Few symptoms” were reported in nursing infants which were considered “not serious.”

## Data Show Widespread Distribution of mRNA

Data shows LNPs can travel throughout the body to various tissues and organs, and therefore logical to think lipid nanoparticles (LNPs) containing mRNA could spread through vaccination to the mammary glands. Byram Bridle, a Canadian virologist and vaccinologist, [obtained a Pfizer biodistribution study](#) in rodents from Japan that showed LNPs could pass through biological tissues and membranes and travel to various organs. In the biodistribution study, 75 percent of LNPs had left the injection site within 48 hours of vaccination and were concentrated in the spleen and liver. Levels were also detected in the ovaries, adrenal glands, brain, eyes, heart, testes, uterus, pituitary gland, spinal cord, thymus, and bone marrow.

Numerous studies have found LNPs containing [mRNA still circulating](#) in the blood after vaccination, with one study ([pdf](#)) showing the presence of mRNA and free spike proteins—for the entire 60-day duration of the study—in the cytoplasm and nuclei of germinal cells in armpit lymph nodes on the same side of the body as the injection site.

A [recently published paper](#) in biomedicines found that LNPs in COVID-19 vaccines containing potentially inflammatory synthetic mRNA do not remain at the injection site after vaccination but are widely distributed throughout the body and can cross protective membranes.

The paper cites a European Medicines Agency report that found, “mRNA could be detected in the brain following intramuscular administration at about 2% of the level found in plasma.” Another study cited in the paper describes how lipid nanoparticles can easily cross the blood-brain barrier.

According to a 2022 study in Biomedicines, [similar vaccine formulations](#) were tracked in various tissues of rodents up to five days post-injection by Moderna and 14 days by Pfizer, but no human biodistribution data are currently available for any of the two licensed products—SPIKEVAX and Comirnaty. In the same study, researchers detected mRNA in blood 15 days following vaccination.


[Other research](#) detected spike proteins in circulating exosomes that carry nucleic acids, proteins, lipids, and metabolites throughout the body for at least four months after vaccination with Pfizer’s COVID-19 vaccine.

\*

Note to readers: Please click the share button above. Follow us on Instagram and Twitter and subscribe to our Telegram Channel. Feel free to repost and share widely Global Research articles.

**Megan Redshaw** is an attorney and investigative journalist with a background in political science. She is also a traditional naturopath with additional certifications in nutrition and exercise science.

Featured image is from [womenshealth.gov](#)



## The Worldwide Corona Crisis, Global Coup d'Etat Against Humanity

by Michel Chossudovsky

Michel Chossudovsky reviews in detail how this insidious project “destroys people’s lives”. He provides a comprehensive analysis of everything you need to know about the “pandemic” — from the medical dimensions to the economic and social repercussions, political underpinnings, and mental and psychological impacts.

*“My objective as an author is to inform people worldwide and refute the official narrative which has been used as a justification to destabilize the economic and social fabric of entire countries, followed by the imposition of the “deadly” COVID-19 “vaccine”. This crisis affects humanity in its entirety: almost 8 billion people. We stand in solidarity with our fellow human beings and our children worldwide. Truth is a powerful instrument.”*

### Reviews

*This is an in-depth resource of great interest if it is the wider perspective you are motivated to understand a little better, the author is very knowledgeable about geopolitics and this comes out in the way Covid is contextualized. —Dr. Mike Yeadon*

*In this war against humanity in which we find ourselves, in this singular, irregular and massive assault against liberty and the goodness of people, Chossudovsky’s book is a rock upon which to sustain our fight. —Dr. Emanuel Garcia*

*In fifteen concise science-based chapters, Michel traces the false covid pandemic, explaining how a PCR test, producing up to 97% proven false positives, combined with a relentless 24/7 fear campaign, was able to create a worldwide panic-laden “plandemic”; that this plandemic would never have been possible without the infamous DNA-modifying Polymerase Chain Reaction test – which to this day is being pushed on a majority of innocent people who have no clue. His conclusions are evidenced by renown scientists. —Peter Koenig*

*Professor Chossudovsky exposes the truth that “there is no causal relationship between the virus and economic variables.” In other words, it was not COVID-19 but, rather, the deliberate implementation of the illogical, scientifically baseless lockdowns that caused the shutdown of the global economy. —David Skripac*

*A reading of Chossudovsky’s book provides a comprehensive lesson in how there is a global coup d’état under way called “The Great Reset” that if not resisted and defeated by freedom loving people everywhere will result in a dystopian future not yet imagined. Pass on this free gift from Professor Chossudovsky before it’s too late. You will not find so much valuable information and analysis in one place. —Edward Curtin*

ISBN: 978-0-9879389-3-0, Year: 2022, PDF Ebook, Pages: 164, 15 Chapters

Price: ~~\$11.50~~ FREE COPY! [Click here \(docsend\)](#) and download.

We encourage you to support the eBook project by making a donation through Global Research’s [DonorBox](#) “**Worldwide Corona Crisis**” [Campaign Page](#).

The original source of this article is [The Epoch Times](#)

## Comment on Global Research Articles on our Facebook page

## Become a Member of Global Research

### Related Articles from our Archives

"How mRNA vaccines could target everything from cancer to the plague": They Will Never Stop, Even If Pfizer & Moderna COVID-19 mRNA Vaccines Were a Complete Failure  
31 March 2023

COVID-19 mRNA Vaccination Campaign Is Mass Homicide  
14 August 2023

International Science Journal Publishes Article Calling for COVID-19 mRNA Vaccines to be Reclassified as Gene Therapies  
7 July 2023



Translate Website

Articles by:

**Megan Redshaw**

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)

[www.globalresearch.ca](http://www.globalresearch.ca) conține materiale protejate prin drepturi de autor a căror utilizare nu a fost întotdeauna autorizată în mod specific de către proprietarul drepturilor de autor. Punem la dispoziția cititorilor noștri un astfel de material în conformitate cu prevederile „utilizării loiale” într-un efort de a promova o mai bună înțelegere a problemelor politice, economice și sociale. Materialul de pe acest site este distribuit fără profit celor care și-au exprimat în prealabil interesul de a-l primi în scop de cercetare și educație. Dacă doriți să utilizați material protejat prin drepturi de autor în alte scopuri decât „utilizarea loială”, trebuie să solicitați permisiunea proprietarului drepturilor de autor.

Pentru întrebări media: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)

Global Research News  
I-BOOKS SERIES  
Countries Index  
Most Popular  
Links  
Contact  
Membership  
Online Store

### Themes

US NATO War Agenda  
Global Economy  
Crimes against Humanity  
Military and WMD  
Law and Justice  
Police State & Civil Rights  
History  
9/11 & 'War on Terrorism'  
Media Disinformation

Militarization and WMD  
Oil and Energy  
Police State & Civil Rights  
Religion  
Poverty & Social Inequality  
Science and Medicine  
United Nations  
US NATO War Agenda  
Women's Rights

### Geographic Regions

[Privacy Policy](#)

Copyright © 2005-2023 GlobalResearch.ca